Subject Index

Aldicarb: Accelerated degradation of aldicarb and its oxidation products in previously treated soils, 147
Alfalfa production: Alfalfa production as influenced by pest stress and early first harvest in Oklahoma, 190
Almond: *see Phytophthora cambivora*, 314
Alternative host plants: *see Heliothis armigera*, 161
Andropogon virginicus: Selective broomsedge (*Andropogon virginicus* L.) control in permanent pastures, 80
Anthonomus grandis: Area-wide suppression of boll weevil (Coleoptera: Curculionidae) populations in Nicaragua, 168
Aphids: *see* Organophosphates, 267
Apple: *see* Miridae, 118
Ascochyta paspali: Survey of the distribution and perceived losses caused by paspalum leaf blight in Northern Victoria, 28
Avena fatua: *see* Barban, 3
Avocado pests: *see Bacillus thuringiensis*, 131

Bacillus thuringiensis: Efficacy of *Bacillus thuringiensis* preparations containing dead and live spores against two avocado pests: the giant looper, *Boarmia selenaria* (Lep.: Geometridae), and the honeydew moth, *Cryptoblabes gnidiella* (Lep.: Phycitidae), 131
Barban: Wild-oat herbicide studies: barban herbicidal activity for wild-oat (*Avena fatua* L.) control as influenced by adjuvants, 3
Bemisia tabaci: *see* Sweet potato whitefly, 43
Bioassay: *see* Prodiamine, 75
Biological control: *see Bacillus thuringiensis*, 131
Biology: *see Euproctis lunata*, 332
Botrytis cinerea: Latent infection of *Botrytis cinerea* in rose flowers and combined chemical and physiological control of the disease, 361
Brassica oleracea: Cabbage development and associated lepidopterous pest complex in the southern USA, 396
Brassica oleracea: Resistance to insects by cabbage lines developed in New York when grown in South Texas, 391
Busseola fusca: Survival of maize stalk borer, *Busseola fusca* (Fuller), in crop residues in Ethiopia, 183

Calcium chloride: *see* Groundnuts, 198
Carbofuran: *see Schizaphis graminum*, 336
Carbosulfan: *see* Cowpea, 384
Carboxin: *see Ustilago nuda*, 16
Cereal crops: *see* Spraying methods, 125
Cereal cultivars: *see* Grasshopper damage, 338
Cereal cyst nematode: *see Heterodera avenae*, 378
Cereals, spring: *see* Spraying methods, 303
Chemical pest control: *see* Spraying methods, 153
Cherry: *see Phytophthora cambivora*, 314
Chrysomphalus aonidum: Effect of malathion–bait mixture on two parasitoids of the Florida red scale, *Chrysomphalus aonidum* (L.), 91
Cocoa: Ghanaian cocoa varieties and swollen shoot virus, 219
Conifers: *see* Organophosphates, 267
Cost–benefit analysis: Cost benefits of integrating nitrogen, plant growth regulators and fungicides in winter barley, 252
Cotton: *see Heliothis armigera*, 161
Cotton: *see* Spraying methods, 153
Cotton: *see* Sweet potato whitefly, 43
Cotton pest management: *see Anthonomus grandis*, 168
Cotton seedling disease control: *see Rhizoctonia solani*, 309
Cowpea: Cowpea seed treatment with carbosulfan: potential for control of seedling pests, 384
Cowpea: Effect of cowpea/maize mixed cropping on the incident light at the cowpea canopy and flower thrips (Thysanoptera: Thripidae) population density, 186
Crop residues: *see Busseola fusca*, 183
Cydia pomonella: *see* Spraying methods, 112

Damage threshold: *see* Miridae, 118
Distribution: *see Ascochyta paspali*, 28
Distribution: *see* Weeds, aquatic, 232

Ecology: *see* Weeds, aquatic, 283
Electrostatically charged sprays: *see* Spraying methods, 125
Euproctis lunata: Biology, seasonal occurrence, host range and damage potential of the castor hairy caterpillar, *Euproctis lunata* Walk. (Lymentriidae: Lepidoptera), 332

Fenfuram: *see Ustilago nuda*, 16
Fenithrothion: *see* Spraying methods, 106
Fluazifop-butyl: *see* Sunflower, 9
Foliar disease: *see* Grasshopper damage, 338
Fungal diseases: *see Triticum aestivum*, 207
Fungicide application methods: *see Rhizoctonia solani*, 309
Fungicides: *see Botrytis cinerea*, 361
Fungicides: Past orchard performance as a guide to decisions on use of fungicides to control sooty blotch and flyspeck, 258
Fungicides: *see* Ryegrass, 34
Fusarium oxysporum: *see Verticillium dahliae*, 327

Gaeumannomyces graminis: *see* Take-all, 20
Gloeodes pomigena: *see* Fungicides, 258
Glycine max: *see* Soybean, 48
Grasshopper damage: Differential feeding by grasshoppers and levels of foliar diseases in various cultivars of spring cereals, 338
Groundnuts: Prevention of condensation damage to groundnuts in containers using calcium chloride, 198

Harvest schedule: *see* Alfalfa production, 190
Heliothis armigera: Significance of host-plant phenology in the dynamics and pest incidence of the cotton bollworm, *Heliothis armigera* Hübner (Lepidoptera: Noctuidae) in Western Tanzania, 161
Herbicide tolerance: *see* Prodiamine, 75
Herbicides: *see Andropogon virginicus*, 80
Herbicides: *see Lathyrus sativus*, 372
Herbicides: *see* Rice seedlings, 202
Heterodera avenae: Observations on the prevalence and incidence of cereal cyst nematode (*Heterodera avenae*) in Scotland, 378
Heterodera cruciferae: *see* Oilseed rape, 137

Insecticide toxicity: *see Nephotettix virescens*, 66
Insecticides: *see* Ryegrass, 34

Lathyrus sativus: Effect of herbicides and annual weeds on the yield and seed quality of lathyrus (*Lathyrus sativus* L.), 372
Leaf production: *see Brassica oleracea*, 396
Leaf rust: Effect of leaf rust (*Puccinia recondita tritici*) on yield of four isogenic wheat lines, 39

Maize: *see* Cowpea, 186
Malathion toxicity: *see Chrysomphalus aonidum*, 91
Metham sodium: *see Sclerotinia sclerotiorum*, 25
Methyl bromide: *see Sclerotinia sclerotiorum*, 25
Miridae: Elements for assessing mirid (Heteroptera: Miridae) damage threshold on apple fruits, 118
Mixed cropping: *see* Cowpea, 186
Myzus persicae: Side-band application of systemic granular pesticides for the control of aphids and potato leafroll virus, 96
Myzus persicae: Insecticide toxicity to three strains of green peach aphid (*Myzus persicae* Sulzer) reared on resistant and susceptible potato cultivars, 62
Myzus persicae: Effect of repeated applications of insecticides to potatoes on numbers of *Myzus persicae* (Sulzer) (Hemiptera: Aphididae) and on the frequencies of insecticide-resistant variants, 55

Nematicides: *see* Oilseed rape, 137
Nephotettix virescens: Comparative toxicity of synthetic pyrethroid, organophosphate and carbamate insecticides against *Nephotettix virescens* (Distant) and *Nilaparvata lugens* (Stål) in rice, 66
Nilaparvata lugens, *see Nephotettix virescens*, 66
Nilaparvata lugens: Operation of a brown planthopper surveillance system in the Tanjung Karang Irrigation Scheme in Malaysia, 273
Nilaparvata lugens: *see* Wild rices, 403
Nitrogen: *see* Cost–benefit analysis, 252

Oilseed rape: Field investigation of the responses to nematicide treatment of three winter oilseed rape cultivars infested by *Heterodera cruciferae*, 137
Onions: *see Stemphylium*, 356
Organophosphates: Comparison of the potential of organophosphorous insecticides and soaps in conifer aphid control, 267
Oryza spp.: *see* Wild rices, 403
Ostrinia nubilalis: Use of strips of *Zea mays* L. as a trap for European corn borer (*Ostrinia nubilalis* Hbn.) oviposition in maize fields, 177

Parasitoids: *see Chrysomphalus aonidum*, 91
Paspalum leaf blight: *see Ascochyta paspali*, 28
Permanent pastures; *see Andropogon virginicus*, 80
Pest complex: *see Brassica oleracea*, 396
Pest management: *see* Rice pests, 84
Pest resistance: *see Brassica oleracea*, 391
Pest stress: *see* Alfalfa production, 190
Pest survival: *see Busseola fusca*, 183
Pesticide residues: *see* Aldicarb, 147
Phenylamides: *see Phytophthora infestans*, 210
Phenylamides: Antifungal modes of action of metalaxyl, cyprofuram, benalaxyl and oxadixyl in phenylamide-sensitive and phenylamide-resistant strains of *Phytophthora megasperma* f.sp *medicaginis* and *Phytophthora infestans*, 347
Phoma betae: Thiram steep seed treatment to control *Phoma betae* and improve the establishment of the sugar-beet plant stand, 319
Phytophthora cambivora: Preliminary evaluation of phosphorous acid, fosetyl-Al and metalaxyl for controlling *Phytophthora cambivora* on almond and cherry, 314
Phytophthora infestans: *see* Phenylamides, 347
Phytophthora infestans: Efficacy over time of cymoxanil in controlling late blight in potatoes incited by a phenylamide-resistant isolate of *Phytophthora infestans*, 210
Phytophthora megasperma: see Phenylamides, 347
Plant growth regulators: *see* Cost–benefit analysis, 252
PLRV: *see Myzus persicae*, 96
Potato: *see Myzus persicae*, 62
Potato: *see Myzus persicae*, 55
Potato: *see Phytophthora infestans*, 210
Potatoes: *see Myzus persicae*, 96
Prodiamine: Comparative tolerance of different plant species to prodiamine, 75
Puccinia recondita tritici: *see* Leaf rust, 39
Puccinia sorghi: Susceptibility of US sweet corn hybrids to *Puccinia sorghi* Schw. in Hawaii, 249

Raspberry: *see* Spraying methods, 106
Resistance: *see Puccinia sorghi*, 249
Resistance breeding: *see Brassica oleracea*, 391
Resistance breeding: *see* Cocoa, 219
Rhizoctonia solani: Evaluation of seed dressing and in-furrow treatments with fungicides for control of seedling disease in cotton caused by *Rhizoctonia solani*, 309
Rhopalosiphum maidis: *see Schizaphis graminum*, 336
Rice: *see Nilaparvata lugens*, 273
Rice pests: An approach to improving pest management: rice in Malaysia, 84
Rice seedlings: Weed control in rice seedling nurseries, 202
Ricinus communis: *see Euproctis lunata*, 332
Rose: *see Botrytis cinerea*, 361
Ryegrass: Improvements to newly sown ryegrass by use of combined fungicide and insecticide treatment, 34

Saudi Arabia: *see Triticum aestivum*, 207
Schizaphis graminum: Effectiveness of carbofuran applied to the soil and as a seed treatment for the control of aphids on wheat in Northwestern Mexico, 336
Sclerotinia sclerotiorum: Control of sclerotia and apothecia of *Sclerotinia sclerotiorum* by metham sodium, methyl bromide and soil solarization, 25
Seedling pest control: *see* Cowpea, 384
Sesame: Manipulation of sowing dates for the control of bacterial blight of sesame (*Sesamum indicum* L.) induced by *Xanthomonas campestris* pv. *sesami* (Sabet & Dowson) Dye, 367
Soil degradation: *see* Aldicarb, 147
Soil solarization: *see Verticillium dahliae*, 327
Soil type: *see Heterodera avenae*, 378
Solanum tuberosum: *see Myzus persicae*, 55, 62, 96
Sowing date: *see* Sesame, 367
Soybean: Evaluation of soybean genotypes for field resistance to stink bugs in Nigeria, 48
Spraying methods: Biological efficiency of electrostatically charged sprays applied by hydraulic nozzles to cereal crops, 125
Spraying methods: Effects of application methods, spray volume rates and pressures on weed control in spring cereals, 303
Spraying methods: Investigations into very-low-volume water-based insecticide application to cotton in Malawi, 153
Spraying methods: Comparison of an air-assisted cross-flow sprayer with a conventional hydraulic sprayer for control of raspberry aphids by fenitrothion, 106
Spraying methods: *see Stemphylium*, 356
Spraying methods: Effects of spraying volume and chemical rate on the control of apple scab (*Venturia inaequalis*) and codling moth (*Cydia pomonella*) in an apple orchard, 112
Stemphylium: Air-assisted placement spraying for *Stemphylium* control in seed onions, 356
Stink bugs: *see* Soybean, 48
Stored-product damage: *see* Groundnuts, 198
Sugar beet: *see Phoma betae*, 319
Sunflower: Grass weed control in sunflower in the spring and kharif (summer) seasons under barani (rainfed) conditions in Pakistan, 9
Surveillance system: *see Nilaparvata lugens*, 273
Sweet corn: *see Puccinia sorghi*, 249
Sweet potato whitefly: *Bemisia tabaci* (Homoptera: Aleyrodidae): populations on different cotton varieties and cotton stickiness in Israel, 43
Swollen shoot virus: *see* Cocoa, 219
Systems analysis: *see* Rice pests, 84

Take-all: Effect of chemical and biological treatments on take-all of winter wheat, 20
Thiram steep: *see Phoma betae*, 319
Trap cropping: *see Anthonomus grandis*, 168
Trap cropping: *see Ostrinia nubilalis*, 177
Triticum aestivum: Fungal diseases of spring wheat in central Saudi Arabia, 207

Ustilago nuda: Resistance to carboxin and fenfuram in *Ustilago nuda* (Jens.) Rostr., the causal agent of barley loose smut, 16

Venturia inaequalis: *see* Spraying methods, 112
Verticillium dahliae: Interaction of soil solarization and metham-sodium in the destruction of *Verticillium dahliae* and *Fusarium oxysporum* f.sp. *vasinfectum*, 327

Weed control: *see* Rice seedlings, 202
Weed control: *see* Spraying methods, 303
Weed management: *see* Weeds, aquatic, 232, 283
Weeds: *see Lathyrus sativus*, 372
Weeds, aquatic: Aquatic weed problems and their management: a review. I. The worldwide scale of the aquatic weed problem, 232
Weeds, aquatic: Aquatic weed problems

and their management: a review. II. Physical control measures, 283
Weeds, grass: *see* Sunflower, 9
Wheat; *see* Leaf rust, 39
Wild oats: *see* Barban, 3
Wild rices: Resistance of wild rices, *Oryza* spp., to the brown planthopper, *Nilaparvata lugens* (Stål) (Homoptera: Delphacidae), 403
Winter wheat: *see* Take-all, 20
Xanthomonas campestris: *see* Sesame, 367
Zea mays: *see Ostrinia nubilalis*, 177
Zygophiala jamaicensis: *see* Fungicides, 258

Author Index

AKPA, A. D., ERINLE, I. D. AND MISARI, S. M. Manipulation of sowing dates for the control of bacterial blight of sesame (*Sesamum indicum* L.) induced by *Xanthomonas campestris* pv. *sesami* (Sabet & Dowson) Dye, 367
AMPOFO, J. K. O. *see* KYAMANYWA, S., 186
ASSEFA GEBRE-AMLAK. Survival of maize stalk borer, *Busseola fusca* (Fuller), in crop residues in Ethiopia, 183
AUGENDRE, M. *see* DERRIDJ, S., 177
AUSTERWELL, M. *see* GRINSTEIN, A., 356

BAUMGÄRTNER, J. *see* SCHAUB, L., 118
BELLO, L. L. *see* JACKAI, L. E. N., 48
BEN-YEPHET, Y. Control of sclerotia and apothecia of *Sclerotinia sclerotiorum* by metham sodium, methyl bromide and soil solarization, 25
BEN-YEPHET, Y., MELERO-VERA, J. M. AND DEVAY, J. E. Interaction of soil solarization and metham-sodium in the destruction of *Verticillium dahliae* and *Fusarium oxysporum* f.sp. *vasinfectum*, 327
BERBERET, R. C. *see* LATHEEF, M. A., 190
BERTHIER, G. *see* LEROUX, P., 16
BOAG, B. *see* BOWEN, S. A., 378
BOAMAH, A. *see* THRESH, J. M., 219
BOND, W. Comparative tolerance of different plant species to prodiamine, 75
BOWEN, S. A., SPAULL, A. M. AND BOAG, B. Observations of the prevalence and incidence of cereal cyst nematode (*Heterodera avenae*) in Scotland, 378
BREWBAKER, J. L. *see* KIM, S. K., 249
BUTLER, JR, G. D., RIMON, D. AND HENNEBERRY, T. J. *Bemisia tabaci* (Homoptera: Aleyrodidae): populations on different cotton varieties and cotton stickiness in Israel, 43

CADDEL, J. L. *see* LATHEEF, M. A., 190
CAMPBELL, M. M., LOVELESS, R. T. AND EVANS, P. T. Effects of spraying volume and chemical rate on the control of apple scab (*Venturia inaequalis*) and codling moth (*Cydia pomonella*) in an apple orchard, 112
CARMI, Y. *see* NAVARRO, S., 198
CHINODYA, R. *see* HILLOCKS, R. J., 309
CHOW, P. N. P. Wild-oat herbicide studies: barban herbicidal activity for wild-oat (*Avana fatua*) L. control as influenced by adjuvants, 3
COHEN, E., PODOLER, H. AND EL-HAMLAUWI, M. Effect of malathion–bait mixture on two parasitoids of the Florida red scale, *Chrysomphalus aonidum* (L.), 91
COHEN, Y. *see* SAMOUCHA, Y., 210
CROWLE, W. L. *see* OLFERT, O., 338

DASHIELL, K. E. *see* JACKAI, L. E. N., 48
DAVIDSE, L. C., GERRITSMA, O. C. M., IDELER, J., PIE, K. AND VELTHUIS, G. C. M. Antifungal modes of action of metalaxyl, cyprofuram, benalaxyl and oxadixyl in phenylamide-sensitive and phenylamide-resistant strains of *Phytophthora megasperma* f.sp *medicaginis* and *Phytophthora infestans*, 347
DAXL, R. G. *see* SWEZEY, S. L., 168
DE HAAN, P. *see* WYSOKI, M., 131
DELUCCHI, V. *see* SCHAUB, L., 118
DERRIDJ, S., LEFER, H., AUGENDRE, M. AND DURAND, Y. Use of strips of *Zea mays* L. as a trap for European corn borer (*Ostrinia nubilalis* Hbn.) oviposition in maize fields, 177
DEVAY, J. E. *see* BEN-YEPHET, Y., 327
DEVONSHIRE, A. L. *see* FFRENCH-CONSTANT, R. H., 55
DICKSON, M. H. *see* EDELSON, J. V., 391
DRYDEN, D. *see* WALL, D. A., 372
DUCZEK, L. *see* OLFERT, O., 338
DURAND, Y. *see* DERRIDJ, S., 177
DURRANT, M. J., PAYNE, P. A., PRINCE, J. F. W. AND FLETCHER, R. Thiram steep seed treatment to control *Phoma betae* and improve the establishment of the sugar-beet plant stand, 319
DZIADYK, D. A. *see* OLFERT, O., 338

EDELSON, J. V., TRUMBLE, J. AND STORY, R. Cabbage development and associated lepidopterous pest complex in the southern USA, 396
EDELSON, J. V. AND DICKSON, M. H. Resistance to insects by cabbage lines developed in New York when grown in South Texas, 391
ELAD, Y. Latent infection of *Botrytis cinerea* in rose flowers and combined chemical and physiological control of the disease, 361
EL-HAMLAUWI, M. *see* COHEN, E., 91
EL-MELEIGI, M. A. Fungal diseases of spring wheat in central Saudi Arabia, 207
ERINLE, I. D. *see* AKPA, A. D., 367
EVANS, K. *see* HARRIS, P. G. W., 137
EVANS, P. T. *see* CAMPBELL, M. M., 112

FFRENCH-CONSTANT, R. H., HARRINGTON, R. AND DEVONSHIRE, A. L. Effect of repeated applications of insecticides to potatoes on numbers of *Myzus persicae* (Sulzer) (Hemiptera: Aphididae) and on the frequencies of insecticide-resistance variants, 55
FLETCHER, R. *see* DURRANT, M. J., 319
FOSTER, G. N. *see* WOODFORD, J. A. T., 96
FRANKEL, H. *see* GRINSTEIN, A., 356
FRIESEN, G. H. *see* WALL, D. A., 372

GERRITSMA, O. C. M. *see* DAVIDSE, L. C., 347
GOLD, H. J. AND SUTTON, T. B. Past orchard performance as a guide to decisions on use of fungicides to control sooty blotch and flyspeck, 258
GORDON, S. C. *see* WOODFORD, J. A. T., 96
GORDON, S. C. AND WILLIAMSON, B. Comparison of an air-assisted cross-flow sprayer with a conventional hydraulic sprayer for control of raspberry aphids by fenitrothion, 106
GRIFFIN, J. L., WATSON, V. H. AND STRACHAN, W. F. Selective broomsedge (*Andropogon virginicus* L.) control in permanent pastures, 80
GRINSTEIN, A., FRANKEL, H., AUSTERWEL, M. AND KRITZMAN, G. Air-assisted placement spraying for *Stemphylium* control in seed onions, 356
GUERRA-SOBREVILLA, L. Effectiveness of carbofuran applied to the soil and as a seed treatment for the control of aphids on wheat in Northwestern Mexico, 356
GUNNER, R. *see* HILLOCKS, R. J., 309

HALL, B. *see* WICKS, T. J., 314
HARRINGTON, R. *see* FFRENCH-CONSTANT, R. H., 55
HARRIS, P. *see* PHILLIPS, M. C., 125
HARRIS, P. G. W. and EVANS, K. Field investigation of the responses to nematicide treatment of three winter oilseed rape cultivars infested by *Heterodera cruciferae*, 137
HENNEBERRY, T. J. *see* BUTLER, JR, G. D., 43

HEONG, K. L. *see* OOI, P. A. C., 273
HEONG, K. L. *see* NORTON, G. A., 84
HILLOCKS, R. J., CHINODYA, R. AND GUNNER, R. Evaluation of seed dressing and in-furrow treatments with fungicides for control of seedling disease in cotton caused by *Rhizoctonia solani*, 309
HINKS, C. F. *see* OLFERT, O., 338

IDELER, J. *see* DAVIDSE, L. C., 347
ISLAM, W., NARGIS, A. AND JOARDER, O. I. Biology, seasonal occurrence, host range and damage potential of the castor hairy caterpillar, *Euproctis lunata* Walk. (Lymentriidae: Lepidoptera), 332
IZHAR, Y. *see* WYSOKI, M., 131

JACKAI, L. E. N., ROBERTS, J. M. F. AND SINGH, S. R. Cowpea seed treatment with carbosulfan: potential for control of seedling pests, 384
JACKAI, L. E. N., DASHIELL, K. E. AND BELLO, L. L. Evaluation of soybean genotypes for field resistance to stink bugs in Nigeria, 48
JOARDER, O. I. *see* ISLAM, W., 332
JORDAN, V. W. L. AND STINCHCOMBE, G. R. Cost benefits of integrating nitrogen, plant growth regulators and fungicides in winter barley, 252
JUKES, A. A. *see* SUETT, D. L., 147

KALODE, M. B. *see* KRISHNAIAH, N. V., 66
KHAN, R. U., MORRISON, I. N. AND RASHID, A. Grass weed control in sunflower in the spring and kharif (summer) seasons under barani (rainfed) conditions in Pakistan, 9
KIM, S. K., BREWBAKER, J. L., LOGRONO, M. AND SRINIVASAN, G. Susceptibility of US sweet corn hybrids to *Puccinia sorghi* Schw. in Hawaii, 249
KRISHNAIAH, N. V. AND KALODE, M. B. Comparative toxicity of synthetic pyrethroid, organophosphate and carbamate insecticides against *Nephotettix virescens* (Distant) and *Nilaparvata lugens* (Stål) in rice, 66
KRITZMAN, G. *see* GRINSTEIN, A., 356
KYAMANYWA, S. AND AMPOFO, J. K. O. Effect of cowpea/maize mixed cropping on the incident light at the cowpea canopy and flower thrips (Thysanoptera: Thripidae) population density, 186

LATHEEF, M. A., CADDEL, J. L., BERBERET, R. C. AND STRITZKE, J. F. Alfalfa production as influenced by pest stress and early first harvest in Oklahoma, 190
LEFER, H. *see* DERRIDJ, S., 177
LEROUX, P. AND BERTHIER, G. Resistance to carboxin and fenfuram in *Ustilago nuda* (Jens.) Rostr., the causal agent of barley loose smut, 16
LEVY, R. S. *see* SAMOUCHA, Y., 210
LEWIS, G. C. Improvements to newly sown ryegrass by use of combined fungicide and insecticide treatment, 34
LOCKWOOD, G. *see* THRESH, J. M., 219
LOGRONO, M. *see* KIM, S. K., 249
LOVELESS, R. T. *see* CAMPBELL, M. M., 112

MELERO-VERA, J. M. *see* BEN-YEPHET, Y., 327
MISARI, S. M. *see* AKPA, A. D., 367
MOODY, K. *see* RAO, A. N., 202
MORRISON, I. N. *see* KHAN, R. U., 9
MURPHY, K. J. Aquatic problems and their management: a review. I. The worldwide scale of the aquatic weed problem, 232
MURPHY, K. J. Aquatic weed problems and their management: a review. II. Physical control measures, 283
MUSAU, D. M. AND PARRY, W. H. Comparison of the potential of organophosphorous insecticides and soaps in conifer aphid control, 267

NARGIS, A. *see* ISLAM, W., 332
NAVARRO, S., PASTER, N. and CARMI, Y. Prevention of condensation damage to groundnuts in containers using calcium chloride, 198
NORDBY, A. *see* SKUTERUD, R., 303
NORTON, G. A. AND HEONG, K. L. An approach to improving pest management: rice in Malaysia, 84
NYAMBO, B. T. Significance of host-plant phenology in the dynamics and pest incidence of the cotton bollworm, *Heliothis armigera* Hübner (Lepidoptera: Noctuidae) in Western Tanzania, 161
NYIRENDA, G. K. C. Investigations into very-low-volume water-based insecticide application to cotton in Malawi, 153

OLFERT, O., HINKS, C. F., WESTCOTT, N. D., CROWLE, W. L., DZIADYK, D. A. AND DUCZEK, L. J. Differential feeding by grasshoppers and levels of foliar diseases in various cultivars of spring cereals, 338
OOI, P. A. C. AND HEONG, K. L. Operation of a brown planthopper surveillance system in the Tanjung Karang Irrigation Scheme in Malaysia, 273
OWUSU, G. K. *see* THRESH, J. M., 219

PARRY, W. H. *see* MUSAU, D. M., 267
PASTER, N. *see* NAVARRO, S., 198
PAVELEY, N. *see* PHILLIPS, M. C., 125
PAYNE, P. A. see DURRANT, M. J., 319
PHILLIPS, M. C., PAVELEY, N. AND HARRIS, P. Biological efficiency of electrostatically charged sprays applied by hydraulic nozzles to cereal crops, 125
PIE, K. *see* DAVIDSE, L. C., 347
PODOLER, H. *see* COHEN, E., 91
PRICE, T. V. *see* WILLIAMS, B. L., 28
PRINCE, J. F. W. *see* DURRANT, M. J., 319

RAMAN, K. V. Insecticide toxicity to three strains of green peach aphid (*Myzus persicae* Sulzer) reared on resistant and susceptible potato cultivars, 62
RAO, A. N. AND MOODY, K. Weed control in rice seedling nurseries, 202
RASHID, A. *see* KHAN, R. U., 9
RIMON, D. *see* BUTLER, JR, G. D., 43
ROBERTS, J. M. F. *see* JACKAI, L. E. N., 384
ROELFS, A. P. *see* SECK, M., 39
ROTHROCK, C. S. Effect of chemical and biological treatments on take-all of winter wheat, 20

SAMOUCHA, Y., LEVY, R. S. AND COHEN, Y. Efficacy over time of cymoxanil mixtures in controlling late blight in potatoes incited by a phenylamide-resistant isolate of *Phytophthora infestans*, 210
SCHAUB, L., STAHEL, W. A., BAUMGÄRTNER, J. AND DELUCCHI, V. Elements for assessing mirid (Heteroptera: Miridae) damage threshold on apple fruits, 118
SECK, M., ROELFS, A. P. AND TENG, P. S. Effect of leaf rust (*Puccinia recondita tritici*) on yield of four isogenic wheat lines, 39
SINGH, S. R. *se* JACKAI, L. E. N., 384
SKUTERUD, R., NORDBY, A. AND TYLDUM, A. Effects of application methods, spray volumes, pressures and herbicide rates on weed control in spring cereals, 303
SPAULL, A. M. *see* BOWEN, S. A., 378
SRINIVASAN, G. *see* KIM, S. K., 249
STAHEL, W. A. *see* SCHAUB, L., 118
STINCHCOMBE, G. R. *see* JORDAN, V. W. L., 252
STORY, R. *see* EDELSON, J. V., 396
STRACHAN, W. F. *see* GRIFFIN, J. L., 80
STRITZKE, J. F. *see* LATHEEF, M. A., 190
SUETT, D. L. AND JUKES, A. A. Accelerated degradation of aldicarb and its oxidation products in previously treated soils, 147
SUTTON, T. B. *see* GOLD, H. J., 258
SWEZEY, S. L. AND DAXL, R. G. Area-wide suppression of boll weevil (Coleoptera: Curculionidae) populations in Nicaragua, 168

TAYLOR, P. A. *see* WILLIAMS, B. L., 28
TENG, P. S. *see* SECK, M., 39
THRESH, J. M., OWUSU, G. K., BOAMAH, A. AND LOCKWOOD, G. Ghanaian cocoa varieties and swollen shoot virus, 219
TRUMBLE, J. *see* EDELSON, J. V., 396
TYLDUM, A. *see* SKUTERUD, R., 303

VELTHUIS, G. C. M. *see* DAVIDSE, L. C., 347
VELUSAMY, R. Resistance of wild rices, *Oryza* spp., to the brown planthopper,

Nilaparvata lugens (Stål) (Homoptera: Delphacidae), 403
WALL, D. A., FRIESEN, G. H. AND DRYDEN, D. Effect of herbicides and annual weeds on the yield and seed quality of lathyrus (*Lathyrus sativus* L.), 372
WATSON, V. H. *see* GRIFFIN, J. L., 80
WESTCOTT, N. D. *see* OLFERT., 338
WICKS, T. J. AND HALL, B. Preliminary evaluation of phosphorous acid, fosetyl-Al and metalaxyl for controlling *Phytophthora cambivora* on almond and cherry, 314
WILLIAMS, B. L., PRICE, T. V. AND TAYLOR, P. A. Survey of the distribution and perceived losses caused by paspalum leaf blight in Northern Victoria, 28
WILLIAMSON, B. *see* GORDON, S. C., 106
WOODFORD, J. A. T., GORDON, S. C. AND FOSTER, G. N. Side-band application of systemic granular pesticides for the control of aphids and potato leafroll virus, 96
WYSOKI, M., DE HAAN, P. AND IZHAR, Y. Efficacy of *Bacillus thuringiensis* preparations containing dead and live spores against two avocado pests: the giant looper, *Boarmia selenaria* (Lep.: Geometridae), and the honeydew moth, *Crytoblabes gnidiella* (Lep: Phycitidae), 131

PLANT SCIENCE TITLES FROM BUTTERWORTHS . . .

Manipulation of Flowering
Edited by **J G Atherton,** University of Nottingham School of Agriculture, UK
1987 448 pages 234 x 165mm Hardcover Illustrated 0 407 00570 6 £65.00

The Production of Grapes and Wines in Cool Climates
David Jackson, Reader in the Department of Horticulture, Landscape and Parks, Lincoln Agricultural College, New Zealand
Danny Schuster
International Winemaking Consultant
1987 192 pages 190 x 255mm Hardcover Illustrated 0 409 78784 1 £40.00

Hormone Action in Plant Development – a critical appraisal
Edited by **G V Hoad, M B Jackson, J R Lenton and R K Atkin.** Long Ashton Research Station, Bristol, UK
1987 336 pages 234 x 156mm Hardcover 0 408 00796 6 £22.00

Temperate and Subtropical Fruit Production
David Jackson, Reader in the Department of Horticulture, Landscape and Parks, Lincoln Agricultural College, New Zealand
1986 302 pages 255 x 190mm Hardcover Illustrated 0 409 70149 1 £40.00 Butterworths, New Zealand
Butterworths, New Zealand

Please order these titles from your usual bookseller – or in case of difficulty or for further information write to Butterworths at the address below.

Butterworths
meeting professional needs

BOROUGH GREEN, SEVENOAKS, KENT, TN15 8PH, UK